1. WFSV Kongress Medizin am Arbeitsplatz

„Betriebliche Gesundheit von Außen“

Dr. Johannes Koch

Facharzt für Innere Medizin

Arzt für Allgemeinmedizin

Gesundheit im Alter

Fit im Alter

Gesund im Arbeitsprozess

BGM

Betriebliches Gesundheitsmanagement

Arbeits- u. Gesundheitsschutz
einschließlich arbeitsmedizinischer Vorsorge

Betriebliches
Eingliederungsmanagement

Betriebliche
Gesundheitsförderung

Partizipation

Interdisziplinarität

Diversität

Digitalisierung

Franklin D. Roosevelt
Präsident der Vereinigten Staaten
von Amerika von 1932 - 1945

1940

- kardiovaskuläre Erkrankungen häufigste Todesursache in den USA
- verantwortlich für jede zweiten Todesfall

Kannel WB. Contribution of the Framingham Study to preventive cardiology. *Journal of the American College of Cardiology*. 1990; 15:206–11. [PubMed: 2136875]

- 1932: RR 140/100 mmHg
- 1940: RR 188/105 mmHg

Blutdruck 188/105 mmHg

„no more than normal for a man his age“

Bungarner, J. The health of Presidents. Jefferson, N.C.: McFarland; 1994.

- 1932: RR 140/100 mmHg
- 1940: RR 188/105 mmHg
- 1944: RR 240/130 mmHg,
Diagnose: Hypertonie,
Herzschwäche, Therapie mit
Phenobarbital, Salzreduktion,
Digitalis

1945: RR 300/190, Tod durch Gehirnblutung

1948

- **National Heart Act**
- „...the Nation health is seriously threatend by diseases of the heart and circulation, including high blood pressure...”

1948

- Framingham Studie
 - epidemiologische Studie

Massachusetts

Framingham Studie: 1948-2017

- Risikofaktoren für Herz-Kreislauf-Erkrankungen
 - Blutdruck
 - Cholesterin
 - Rauchen
 - Übergewicht
- Konzept der Prävention
- Grenzwerte für Blutdruck und Cholesterin definiert

Table 5 Comparison of office, home, and ambulatory blood pressure measurement thresholds for elevated blood pressure and hypertension

	Office BP (mmHg) ^a	Home BP (mmHg)	Daytime ABPM (mmHg)	24 h ABPM (mmHg)	Night-time ABPM (mmHg)
Reference					
Non-elevated BP	<120/70	<120/70	<120/70	<115/65	<110/60
Elevated BP	120/70–<140/90	120/70–<135/85	120/70–<135/85	115/65–<130/80	110/60–<120/70
Hypertension	≥140/90	≥135/85	≥135/85	≥130/80	≥120/70

© ESC 2024

ABPM, ambulatory blood pressure monitoring; BP, blood pressure.

^aThe BP thresholds provided assume that a standardized approach to office BP measurement is performed (Figure 3). However, evidence indicates that office BP measurement in routine clinical settings is often not done using a standardized approach and, in this case, the routine office BP value may be 5–10 mmHg higher than it would have been if measured using the recommended standardized approach.^{65,66}

Richtlinien 2024

ESC

European Society
of Cardiology

European Heart Journal (2024) **45**, 3912–4018

<https://doi.org/10.1093/eurheartj/ehae178>

ESC GUIDELINES

2024 ESC Guidelines for the management of elevated blood pressure and hypertension

< 40 Jahre – Blutdruckmessung alle 3 Jahre

> 40 Jahre – Blutdruckmessung einmal jährlich

Office blood pressure measurement

1
Measure after 5 min seated comfortably in a quiet environment

2
Use a validated device with an appropriate cuff size based on arm circumference

3
Place the BP cuff at the level of the heart with the patient's back and arm supported

8
Assess for orthostatic hypotension at 1st visit and thereafter by symptoms

4
Measure BP three times (1–2 min apart) and average the last 2 readings

7
Record heart rate and exclude arrhythmia by pulse palpation

6
Measure BP in both arms at the 1st visit to detect between arm differences

5
Obtain further measurements if the readings differ by >10 mmHg

Rauchen
hohes Cholesterin
hoher Blutdruck

Arteriosklerose =
Gefäßverkalkung

Herzinfarkt

Wie hoch darf mein Cholesterinspiegel sein?

Ab wann muss ich Medikamente für einen erhöhten Cholesterinspiegel nehmen?

Examination date:

16 September 2025

Age: 40 (7/1985)

Sex: male

Systolic blood pressure:
125 mmHg

Total Cholesterol: 220 mg/dl

HDL-Cholesterol: 45 mg/dl

LDL-Cholesterol: 130 mg/dl

Current Smoker: No

Examination date:

16 September 2025

Age: 65 (7/1960)

Sex: male

Systolic blood pressure:
125 mmHg

Total Cholesterol: 220 mg/dl

HDL-Cholesterol: 45 mg/dl

LDL-Cholesterol: 130 mg/dl

Current Smoker: No

Your 10-year risk of fatal and non-fatal CVD events is * :

2.2%

For a person of your age, the Guidelines consider a 10-year CVD risk to be high if above:

2.5%

Your 10-year risk of fatal and non-fatal CVD events is * :

9.1%

For a person of your age, the Guidelines consider a 10-year CVD risk to be high if above:

5%

Examination date:

16 September 2025

Age: 65 (7/1960)

Sex: male

Systolic blood pressure:
125 mmHg

Total Cholesterol: 220 mg/dl

HDL-Cholesterol: 45 mg/dl

LDL-Cholesterol: 130 mg/dl

Current Smoker: No

Examination date:

16 September 2025

Age: 65 (7/1960)

Sex: male

Systolic blood pressure:
125 mmHg

Total Cholesterol: 220 mg/dl

HDL-Cholesterol: 45 mg/dl

LDL-Cholesterol: 130 mg/dl

Current Smoker: Yes

Your 10-year risk of fatal and non-fatal CVD events is * :

9.1%

For a person of your age, the Guidelines consider a 10-year CVD risk to be high if above:

5%

Your 10-year risk of fatal and non-fatal CVD events is * :

14%

For a person of your age, the Guidelines consider a 10-year CVD risk to be high if above:

5%

Was Faktoren hängt
die Höhe meines
Cholesterinspiegels
ab?

- Gene
- Bewegung
- Ernährung (gesättigten Fettsäuren, Transfette)

Abbildung 4.1:

Anteil der täglich Rauchenden an der Bevölkerung 15+ 1972-2022

Sind E-Zigaretten gesünder?

- enthalten weniger Schadstoffe
- Nikotin selber ist nicht so schädlich
- aber andere Aerosole und Inhaltsstoffe – Langzeitauswirkungen?
- Abhängigkeitspotential
- deswegen nicht empfohlen
- evtl. Einsatz in der Entwöhnung

- Chronische obstruktive Lungenerkrankung - COPD

Sitzen ist das neue Rauchen!

„Couchpotato“

Zivilisationskrankheiten

Übergewicht und Krankheiten

Figure 1. Effect of Five Modifiable Cardiovascular Risk Factors on Lifetime Risk of Cardiovascular Disease and Death from Any Cause.

Shown are risk curves for participants without the five cardiovascular risk factors (solid lines) as compared with those with all five risk factors (dashed lines) at an index age of 50 years. Cumulative incidence curves (Panels A and B) and survival curves (Panels C and D) are shown for women (red) and men (blue). Lifetime risk is shown for cardiovascular disease (Panel A) and death from any cause (Panel B), and lifetime difference is shown for cardiovascular disease or death (Panel C) and death from any cause (Panel D). In Panels C and D, cardiovascular disease-free life expectancy and overall life expectancy are indicated by the age at which a given survival curve crosses the horizontal line at 50%. The curves were generated with the use of recalibrated predictions from Weibull models. The five risk factors are a systolic blood pressure of 130 mm Hg or more, a non-high-density lipoprotein (HDL) cholesterol level of 130 mg per deciliter (3.36 mmol per liter) or more, a body-mass index (BMI) of less than 20 or 25 or more, diabetes, and current smoking.

Blutdruck > 130/- mmHg
 Cholesterin
 Übergewicht oder
 Untergewicht
 Diabetes
 Rauchen

Verlust von 10
 Lebensjahren

BGM

Betriebliches Gesundheitsmanagement

Arbeits- u. Gesundheitsschutz
einschließlich arbeitsmedizinischer Vorsorge

Betriebliches
Eingliederungsmanagement

Betriebliche
Gesundheitsförderung

Partizipation

Interdisziplinarität

Diversität

Digitalisierung

betriebliche Gesundheit – internistische Sicht

- Animation zur Bewegung (Lift, Weg zur Arbeit nutzen, Fitnessraum, gemeinsame Aktivitäten...)

betriebliche Gesundheit – internistische Sicht

- Animation zur Bewegung (Lift, Weg zur Arbeit, Fitnessraum, Kooperation mit Fitnesscenter, gemeinsame Aktivitäten...)
- Gesundes Essen

betriebliche Gesundheit – internistische Sicht

- Animation zur Bewegung (Lift, Weg zur Arbeit, Fitnessraum, Kooperation mit Fitnesscenter, gemeinsame Aktivitäten...)
- Gesundes Essen
- Bereitstellung niederschwelliger med. Beratung und Vorsorge (Gesundenuntersuchungen)

Gesundheit im Alter

High risk	<p>People with any of the following:</p> <ul style="list-style-type: none">• Markedly elevated single risk factors, in particular TC >8 mmol/L (>310 mg/dL), LDL-C >4.9 mmol/L (>190 mg/dL), or BP $\geq$180/110 mmHg• Patients with FH without other major risk factors• Patients with DM without target organ damage,^c with DM duration $\geq$10 years or another additional risk factor• Moderate CKD (eGFR 30–59 mL/min/1.73 m²)• A calculated SCORE2 or SCORE2-OP $\geq$10% and <20% for 10 year risk of fatal or non-fatal CVD
Moderate risk	<p>People with any of the following:</p> <ul style="list-style-type: none">• Young patients (T1DM <35 years; T2DM <50 years) with DM duration <10 years, without other risk factors• Calculated SCORE2 or SCORE2-OP $\geq$2% and <10% for 10 year risk of fatal or non-fatal CVD
Low risk	<ul style="list-style-type: none">• Calculated SCORE2 or SCORE2-OP <2% for 10 year risk of fatal or non-fatal CVD

Table 4 Intervention strategies as a function of total cardiovascular risk and untreated low-density lipoprotein cholesterol levels

Total CV risk	Untreated LDL-C levels					
	<1.4 mmol/L (<55 mg/dL)	1.4 to <1.8 mmol/L (55 to <70 mg/dL)	1.8 to <2.6 mmol/L (70 to <100 mg/dL)	2.6 to <3.0 mmol/L (100 to <116 mg/dL)	3.0 to <4.9 mmol/L (116 to <190 mg/dL)	≥4.9 mmol/L (≥190 mg/dL) ^a
Low	Lifestyle advice	Lifestyle advice	Lifestyle advice	Lifestyle advice	Lifestyle modification, consider adding drug if uncontrolled	N/A ^a
Moderate	Lifestyle advice	Lifestyle advice	Lifestyle advice	Lifestyle modification, consider adding drug if uncontrolled	Lifestyle modification, consider adding drug if uncontrolled	N/A ^a
High	Lifestyle advice	Lifestyle advice	Lifestyle modification, consider adding drug if uncontrolled	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention
Very high: primary prevention	Lifestyle modification, consider adding drug	Lifestyle modification, consider adding drug	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention
Very high: secondary prevention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention	Lifestyle modification and concomitant drug intervention